

地球惑星科学専攻 地球科学輻合部特別講演会

2009年5月1日（金曜日）16時30分～

京都大学理学部2号館1階第3講義室

Rheology of serpentines, seismicity and mass transfer in subduction zone

Bruno Reynard

ERI, Univ. Tokyo; Ecole Normale Supérieure de Lyon

講演者紹介：

Serpentinized rocks have a lower density and lower viscosity than "dry" ultramafic rocks and it was proposed, based on numerical simulations, that they play a major role in mantle-slab decoupling, and in downward (sink) or upward (exhumation) motion of eclogites and ultra-high pressure (UHP) rocks in subduction zones. Rheological data on antigorite, the stable variety of serpentine in subduction zones, were obtained over a P-T range of 1-4 GPa and 200-500 /deg C that cover most of its stability field. The experiments were carried out in a D-DIA apparatus installed at GSECARS on the 13-BM-D line of APS. The determined stress-strain curves were fitted to a power-law equation including both temperature and pressure dependence. The results confirm that serpentinites acts as a weak layer that allows significant mass transfer along the "serpentinized channel" and dynamic processes such as mantle slab decoupling, and mantle wedge convection. Regardless of the temperature, serpentinitized mantle at the slab surface has a low viscosity that allows localizing the deformation and impeding stress build-up. It will limit the downdip propagation of large earthquakes, and allow viscous relaxation as an origin of post-seismic deformations and slow earthquakes. Models of growth and transport of a serpentinitized channel using available kinetic and present rheological data explain high exhumation rates of eclogites and limited thickness of the channel at great depths (? 50 km), and slower exhumation in a thick hydrated mantle corner at shallower depths. Such channels may be difficult to detect from seismic tomography or using guided waves because of their small thickness (<2-3 km).

問い合わせ先：川本 竜彦